

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number
WO 01/90130 A1

- (51) International Patent Classification?: C07H 19/16, C12P 19/40, C07H 1/08
- (21) International Application Number: PCT/EP01/03633
- (22) International Filing Date: 30 March 2001 (30.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
MI2000A001158 25 May 2000 (25.05.2000) IT
- (71) Applicant (for all designated States except US): CHE-
MENTECNO S.R.L. [IT/IT]; Via Catalani, 5, I-20052
Monza (IT).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): BERNA, Marco
[IT/IT]; Via Catalani, 5, I-20052 Monza (IT). SIVIERI,
Lino [IT/IT]; Via Catalani, 5, I-20052 Monza (IT). SAN-
TAMBROGIO, Gianni [IT/IT]; Via Catalani, 5, I-20052
Monza (IT). VALOTI, Ermanno [IT/IT]; Via Catalani, 5,
I-20052 Monza (IT).
- (74) Agents: MANCINI, Vincenzo et al.; Ing. A. Giambro-
cono & C. s.r.l., Via Rosolino Pilo, 19/b, I-20129 Milan
(IT).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report
— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments
- For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 01/90130 A1

(54) Title: PROCESS FOR THE PREPARATION OF PHARMACEUTICALLY ACCEPTABLE SALTS OF (SS-RS)-S-ADENO-
SYL-L-METHIONINE

(57) Abstract: The present invention relates to a process for the preparation of pharmaceutically acceptable salts of (SS, RS)-S-adenosyl-L-methionine and allows to obtain the salified (RS)-(+)-S-adenosyl-L-methionine diastereoisomer in amounts lower than or equal to 3 % with respect to the salified (SS)-(+)-S-adenosyl-L-methionine diastereoisomer; the salts that can be obtained by the process of the invention keep their configuration stable in time.


**PROCESS FOR THE PREPARATION OF PHARMACEUTICALLY ACCEPTABLE
SALTS OF (SS,RS)-S-ADENOSYL-L-METHIONINE**

The present invention relates to a process for the
5 preparation of pharmaceutically acceptable salts of
(SS,RS)-S-adenosyl-L-methionine (hereinafter referred to
as (SS,RS)-SAME).


In particular, the invention relates to a process for
the preparation of pharmaceutically acceptable salts of
10 (SS,RS)-SAME, wherein the salified (RS)-(+)-S-adenosyl-L-
methionine diastereoisomer (hereinafter referred to as
(RS)-(+)-SAME) is produced in amounts lower than or equal
to 3% with respect to the salified (SS)-(+)-S-adenosyl-L-
methionine diastereoisomer (hereinafter referred to as
15 (SS)-(+)-SAME).

As it is known, (SS,RS)-SAME is a physiological methyl
donor involved in enzymatic transmethylation reactions,
that is present in all living organisms and has
therapeutical effects on chronic hepatic diseases,
20 adiposis, lipaemia, atherosclerosis and it is desirable,
therefore, to produce it in high amounts.

It is also known, (J. W. Cornforth, J.A.C.S., 1977, 99,
7292-7300; Stolowitz et al., J.A.C.S., 1981, 103, 6015-
6019) that the products containing (SS,RS)-SAME consist of
25 a mixture of two diastereoisomers: (RS)-(+)-SAME and (SS)-
(+)-SAME, having the following structural formulae:



SS



RS

- 2 -

Moreover, it was demonstrated (De La Haba et al., J.A.C.S., 1959, 81, 3975-3980) that only one of the two diastereoisomers, i.e. (SS)-(+)-SAME, is enzymatically active for the transmethylation and spontaneously racemises, thereby giving rise to the formation of the inactive diastereoisomer (RS)-(+)-SAME in a percentage equal to about 20% (Wu et al., Biochemistry 1983, 22, 2828-2832).

The Applicant, in fact, has noted that in all the commercially available products based on (SS,RS)-SAME, the inactive diastereoisomer (RS)-(+)-SAME is present in percentages equal to at least 20%; it was also noted that said percentages increase in time even up to 40% and more.

This observation clearly confirms that the diastereoisomer mixture is unstable in time, which, on the other side, had already been noted in relation with the product in solution (G. L. Creason et al., Phytochemistry, vol. 24, N. 6, 1151-1155, 1985; H. C. Uzar, Liebigs Ann. Chem. 1989, 607-610).

The demand for (SS,RS)-SAME derivatives wherein the percentage of the active (SS)-(+)-SAME diastereoisomer is clearly higher with respect to the inactive (RS)-(+)-SAME isomer and wherein said percentage turns out to be stable in time, is particularly felt in the field.

It was also found that there is an obstacle to the use of (SS,RS)-SAME and the pharmaceutically acceptable salts thereof at the industrial level because of their thermal instability, even at room temperature, and of the complexity of the preparation and purification processes thereof.

Several processes for the purification of (SS,RS)-SAME and for the production of the pharmaceutically acceptable salts thereof are known.

However, the known purification processes, besides providing the use of strong acid resins (JP 13680/1971) or

chelate-type resins (JP 20998/1978) or particular and expensive reactants, such as picric or picolinic acid (US 3707536 and US 3954726), bring anyhow to the partial racemisation of the sulphur chiral center of (SS,RS)-SAME and, therefore, lead to final products containing the inactive diastereoisomer in amounts higher than 20%.

Purification processes that use weak acid resins are also known (JP 14299/1981, FR-A-2531714, EP-A-0141914), which allow, however, to obtain just a partial separation of (SS,RS)-SAME and, therefore, an insufficient purity degree for pharmaceutical purposes.

Even if the realization of some of the above-identified processes enables to obtain a higher purity, the partial racemization implies, at any event, that at least 20% of the inactive diastereoisomer should be present; in some cases moreover (FR-2531714), in order to extract the product from the cells, there is provided the use of potassium bicarbonate, with subsequent precipitation of potassium perchlorate, which brings about problems firstly in the separation and then in the disposal of the product. In EP-A-0141914, the lysis of the cells of the yeast containing (SS,RS)-SAME is carried out in the presence of an organic solvent (for example, ethyl acetate, acetone, etc.) by using, moreover, chromatographic columns based on 100-200 mesh resins, with high investment and maintaining costs. The use of solvents for the extraction of (SS,RS)-SAME necessarily implies the employment of antideflagrant plants and a recovery, distillation and solvent recovery systems, besides the necessary drying of the exhausted mycelium, in order to avoid that it is discharged with the residual solvent, all these factors clearly bringing about additional investment and operation costs.

According to a first aspect, the present invention relates to a process for the preparation of

pharmaceutically acceptable salts of (SS,RS)-SAmE, wherein the salified (RS)-(+)-SAmE diastereoisomer is present in amounts lower than or equal to 3% with respect to the salified (SS)-(+)-SAmE diastereoisomer, which, at a temperature higher than or equal to 0-12°C, comprises:

- the purification of (SS,RS)-S-adenosyl-L-methionine from enriched yeast, which shall contain at least 6 g/l thereof, which comprises:

- (a) - the adjustment of the pH value to 1.2-3.5;
- 10 (b) - the preparation of an aqueous lysate of (SS,RS)-SAmE from the enriched yeast;
- (c) - the microfiltration of the resulting lysate;
- (d) - the absorption of the resulting microfiltrate on a weak acid resin, by eluting with a 0.1-2 N inorganic acid solution;
- 15 (e) - the decolouration of the resulting eluate;
- the concentration of the decolourised eluate, by reverse osmosis, from 30 to 70% by volume;
- the addition of stoichiometric amounts of at least one pharmaceutically acceptable acid salt to the concentrated eluate, so as to obtain the corresponding pharmaceutically acceptable salt of (SS,RS)-SAmE.
- 20

According to a preferred aspect, the so obtained pharmaceutically acceptable salt of (SS,RS)-SAmE can be subjected to lyophilization.

According to another preferred aspect, the process of the invention is carried out at a temperature of 2-5°C.

According to a further preferred aspect, the pH value in step (a) is 1-2, whereas the preparation of the lysate in step (b) can take place by passing the yeast through a breaking-cells equipment, then proceeding with the microfiltration of the so obtained yeast, for example on a ceramic membrane.

The enriched yeast on which (SS,RS)-SAmE is purified preferably contains at least 8-10 g/l of (SS,RS)-SAmE; the

pharmaceutically acceptable acid is selected, preferably, from sulphuric acid and paratoluensulphonic acid.

It can be noted that the process of the invention allows to use resin/product ratios equal to, for example, 5 10-20 liters of resin pro kg of absorbed product, which are advantageous with respect to what has been disclosed in JP 20998/1978.

The process of the invention allows to produce salts of (SS,RS)-SAME wherein, even at room temperature, it is 10 possible to detect a percentage of the (SS)-(+)-SAME diastereoisomer equal to at least 97 with respect to the (RS)-(+)-SAME diastereoisomer which is present, accordingly, in percentages lower than or equal to 3.

The process of the invention allows moreover to 15 exclude the use of organic solvents in the preparation of the lysate, with remarkable advantages with respect to the purification steps of the pharmaceutically acceptable salts of (SS,RS)-SAME, as well as ecological and environmental advantages.

It is furthermore possible to obtain a higher yield 20 and purity of the pharmaceutically acceptable salts of (SS,RS)-SAME with respect to those obtainable by known processes; a purity equal to at least 98% in (SS,RS)-SAME and a yield equal to at least 90 are obtained, in fact, 25 with respect to the fermented product.

Thanks to its particular conditions, the process of the invention allows to avoid the degradation of (SS,RS)-SAME during the preparation of the lysate and allows to obtain a lysis with a yield higher than 98% and with a 30 content of by-products, the main product of which being 5-deacyl-5-methylthioadenosine, lower than 1%.

(SS,RS)-SAME, suitably salified as above described, can be produced, for example, by fermenting a suitable microorganism, such as *Saccharomyces pastorianus* (ex 35 *Saccharomyces carlsbergensis* CBS 1513), *Saccharomyces*

cerevisiae (IFO 2044), *Torulopsis utilis* and *Candida utilis*.

The yeast containing (SS,RS)-SAME can be enriched by the processes known in the field, such as for example, the
5 Schlenk method described in "Journal of Biological Chemistry", vol. 29, page 1037, (1987), which was modified only in optimizing the use of DL methionine and which was conducted at a maximum temperature of 27,5°C for about 20 hours.

10 The (SS,RS)-SAME-enriched yeast (which, in order to be advantageously employed in the realization of the present invention, indicatively contains at least 6 g/l of (SS,RS)-SAME, undergoes, upon adjustment of the pH value to 1.2-3.5, a cellular lysis process, by passing the
15 yeast, preferably, through a cell-breaking equipment.

The resulting lysate, after being subjected to microfiltration, for example on a ceramic membrane such as Kerasep® K09A, is adsorbed on a weak acid carboxylic resin, preferably of the cationic type, such as Rohm and
20 Haas® IRC86, preferably until saturation (about 150 g/l), and eluted with a solution of an inorganic acid such as, for example, 0.1-2 N sulphuric acid, hydrochloric acid, etc.

The decolouration of the resulting eluate takes then
25 place, for example by means of a copolymer resin with a styrene-divinylbenzene unit, such as Resindion® 825L.

The resulting eluate containing (SS,RS)-SAME is concentrated, by reverse osmosis, from 30 to 70%, preferably from 40 to 50% by volume. The so obtained
30 concentrate is added with stoichiometrically equivalent amounts of an acid or a mixture of pharmaceutically acceptable acids, such as those indicated above. The so obtained products can be used for possible preparations in solution or can be subjected to lyophilisation, when one

- 7 -

wishes to use them in the solid form.

The following examples illustrate the invention without limiting it.

EXAMPLE 1

5 1000 kg of yeast obtained by fermentation of
 Saccharomyces carlsbergensis were enriched with (SS,RS)-
 SAME according to the Schlenk method, modified as follows.
 The yeast was added with 100 kg of yeast cream (which,
 upon dilution with 100 l of deionised water has a 2.2 g/l
10 titer), 2 kg of DL methionine, 12 kg of hydrated glucose
 and 1.5 kg of citric acid, keeping under stirring at 27°C
 ± 0,5°C for 22 hours, aerating through emission of sterile
 filtered air at a flow of 0.6 l/l/m, thereby obtaining 9
 g/l of (SS,RS)-SAME. After adjusting pH at 1.2 by means of
15 H₂SO₄, lysis was carried out, at a temperature of 12°C, by
 the "Constant Cell Disruption System" produced by Constant
 System Ltd., a pressure-type cell-breaking system with a
 cooling system. The solution was then cooled by using
 first cold water and then brine, until the solution was
20 brought to a temperature of about 2°C.

 The obtained mixture was then conveyed to a
 microfiltration plant, endowed with cartridges of the type
 Verind A-10 HFM 180 SM, for separating the exhausted solid
 from the enriched liquid. The panel was washed with 2000 l
25 of demineralised water at 2°C. The filtration yield was
 98%.

 The enriched solution was passed through the IRC 86
 resin (Rohm and Haas®), a carboxylic resin and eluted with
 1 N sulphuric acid, still keeping the temperature at about
30 2°C.

 The collected eluate was decolorised by using a
 Resindion® 825L resin. The enriched solution was
 concentrated by reverse osmosis until a 40% concentration
 of (SS,RS)-SAME was obtained. Corresponding stoichiometric
35 amounts of sulphuric acid and paratoluensulphonic acid

were then added to give the disulphate paratoluensulphonate of (SS,RS)-SAME. The final yield of (SS,RS)-SAME disulphate paratoluensulphate was 90%.

The content of (RS)-(+)-SAME disulphate paratoluensulphonate in the diastereoisomer mixture of (SS,RS)-SAME disulphate paratoluensulphonate, analyzed by HPLC, turned out to be 1%. The relevant data are reported in the following, in the table concerning sample N. 4.

EXAMPLE 2

10 1000 kg of yeast, obtained by fermentation of *Saccharomyces carlsbergensis* enriched with (SS,RS)-SAME according to the method described in EXAMPLE 1, with an activity equal to 8.2 g/kg, were lysated by a cell-breaking system at a temperature of 12°C and at a 2 pH.
15 After adding 500 l of water to the resulting solution, the microfiltration and the subsequent steps were carried out, analogously to what described in EXAMPLE 1, washing with 2000 l of cold demineralized water (about 5°C). 7,5 kA of (SS,RS)-SAME were obtained which, after being concentrated
20 by reverse osmosis, were salified obtaining a 91,4% yield of (SS,RS)-SAME disulphate paratoluensulphonate (lysis yield: 99%; purification yield: 98%). The time elapsing from the end of the fermentation to the concentration by reverse osmosis was 32 hours. The relevant data are
25 reported in the following, in the table concerning sample N. 5.

EXAMPLE 3

The solution obtained by the process of EXAMPLE 1, after absorption on IRC 86 (Rohm and Haas®) resin, was
30 eluted with 1 N sulphuric acid.

The obtained solution was concentrated up to 20% and then added with sulphuric acid and paratoluensulphonic acid in a stoichiometric amount, thereafter it was further concentrated until a 40% solution was obtained. 14.09 kg

- 9 -

of (SS,RS)-SAmE disulphate paratoluensulphonate were obtained, with a transformation yield of 97.8%. The relevant data are reported in the following, in the table relating to sample N. 6.

5 EXAMPLE 4 (comparative)

3 samples of SAMIR® [(SS,RS)-SAmE], produced by Knoll Farmaceutici S.p.A., were analyzed by HPLC. The values measured for each sample are as follows:

10 SAMPLE 1 - 100 mg of SAMIR® (vials) batch 045-021;
expiration date 06/2000.

peak No.	retention time	peak area	peak height	area (%)	height (%)
1	3.661	0.26322	0.00171	0.286	0.410
2	4.246	0.33608	0.00210	0.365	0.503
3	4.591	1.82467	0.00906	1.984	2.166
4	5.429	1.00324	0.00573	1.090	1.370
5	5.888	51.25485	0.25301	55.715	60.500
6	6.206	37.31255	0.14658	40.560	35.051

Peak No. 5, corresponding to (SS)-(+)-SAmE, indicates a percentage of 58%, whereas peak No. 6, corresponding to (RS)-(+)-SAmE, indicates a percentage of 42%.

15 SAMPLE 2 - 200 mg of SAMIR® (tablets); batch 121;
expiration date 05/2002.

peak No.	retention time	peak area	peak height	area (%)	height (%)
1	3.655	0.35979	0.00221	0.194	0.269
2	4.238	0.40764	0.00265	0.220	0.322
3	4.538	3.58281	0.01624	1.932	1.973
4	5.411	1.60136	0.00919	0.863	1.116
5	5.828	108.11943	0.52553	58.299	63.833
6	6.144	71.38583	0.26746	38.492	32.487

Peak No. 5, corresponding to (SS)-(+)-SAmE, indicates a percentage of 60%, whereas peak No. 6, corresponding to

(RS)-(+) -SAmE, indicates a percentage of 40%.

SAMPLE 3 - 400 mg of SAMIR® (tablets); batch 040;
expiration date 10/2002.

Peak No.	retention time	peak area	peak height	area (%)	height (%)
1	3.386	0.03675	0.00041	0.010	0.027
2	5.419	1.40489	0.00853	0.387	0.559
3	5.785	214.15843	0.99534	58.973	65.233
4	6.092	147.47305	0.52125	40.610	34.162
5	13.468	0.07286	0.00029	0.020	0.019

Peak No. 3, corresponding to (SS)-(+) -SAmE, indicates a percentage of 59%, whereas peak No. 4, corresponding to (RS)-(+) -SAmE, indicates a percentage of 41%.

EXAMPLE 5

The products obtained according to the process of the invention in EXAMPLES 1-3, samples 4-6 respectively, were analyzed, similarly to what has been described in example 4, after four months from the date of their production.

The measured values for each sample were as follows:

SAMPLE 4 (EXAMPLE 1); batch 003/R.

peak No.	retention time	peak area	peak height	area (%)	height (%)
1	2.595	10.89547	0.06085	4.340	4.566
2	2.735	7.93823	0.07825	3.163	5.873
3	2.834	8.13165	0.08741	3.239	6.561
4	2.946	20.91077	0.12978	8.331	9.740
5	3.355	5.91998	0.02933	2.358	2.201
6	3.651	1.91541	0.00909	0.763	0.683
7	4.136	192.60315	0.92893	76.728	69.716
8	4.958	1.81995	0.00603	0.725	0.453
9	6.423	0.88589	0.00276	0.353	0.207

Peak No. 7, corresponding to (SS)-(+) -SAmE, indicates a percentage of 99%, whereas peak No. 8, corresponding to (RS)-(+) -SAmE, indicates a percentage of 1%.

SAMPLE 5 (Example 2); KF = 2.3%; titre = 102.6%; batch 001/R.

peak No.	retention time	peak area	peak height	area (%)	height (%)
1	2.588	0.23402	0.00223	0.075	0.155
2	2.817	0.08934	0.00099	0.029	0.068
3	2.908	0.28759	0.00228	0.092	0.158
4	3.082	6.84701	0.04649	2.194	3.227
5	3.388	0.69924	0.00391	0.224	0.272
6	3.697	0.84270	0.00466	0.270	0.323
7	4.224	295.93649	1.35851	94.844	94.306
8	5.153	5.50257	0.01712	1.763	1.188
9	6.696	1.58749	0.00436	0.509	0.303

Peak No. 7, corresponding to (SS)-(+)-SAME, indicates a percentage of 98%, whereas peak No. 8, corresponding to (RS)-(+)-SAME, indicates a percentage of 2%.

SAMPLE 6 (EXAMPLE 3); KF = 1.39%; titre = 102.7; batch 004/R.

peak No.	retention time	peak area	peak height	area (%)	height (%)
1	2.584	0.19250	0.00169	0.058	0.109
2	2.825	0.13395	0.00135	0.041	0.088
3	2.894	0.22074	0.00196	0.067	0.127
4	3.060	6.91884	0.04741	2.094	3.072
5	3.355	0.82868	0.00484	0.251	0.314
6	3.661	1.53681	0.00907	0.465	0.588
7	4.162	313.00031	1.45354	94.736	94.209
8	5.026	5.89462	0.01854	1.784	1.201
9	6.528	1.66553	0.00450	0.504	0.292

Peak No. 7, corresponding to (SS)-(+)-SAME, indicates a percentage of 98%, whereas peak No. 6, corresponding to (RS)-(+)-SAME, indicates a percentage of 2%.

CLAIMS

1. A process for the preparation of pharmaceutically acceptable salts of (SS,RS)-SAmE, wherein the salified (RS)-(+)-SAmE diastereoisomer is present in amounts lower
5 than or equal to 3% with respect to the salified (SS)-(+)-SAmE diastereoisomer, which, at a temperature of 0-12°C, comprises:
- the purification of (SS,RS)-S-adenosyl-L-methionine from enriched yeast, which shall contain at least 6 g/l
10 thereof, which comprises:
 - (a) - the adjustment of the pH value to 1.2-3.5;
 - (b) - the preparation of an aqueous lysate of (SS,RS)-SAmE from the enriched yeast;
 - (c) - the microfiltration of the resulting lysate;
 - 15 (d) - the absorption of the resulting microfiltrate on a weak acid resin, by eluting with a 0.1-2 N inorganic acid solution;
 - (e) - the decolouration of the resulting eluate;
 - the concentration of the decolourised eluate by
20 reverse osmosis, from 30 to 70% by volume;
 - the addition of stoichiometric amounts of at least a pharmaceutically acceptable acid to the concentrated eluate, so as to obtain the corresponding pharmaceutically acceptable salt of (SS,RS)-SAmE.
- 25 2. Process according to claim 1, wherein the pharmaceutically acceptable salt of (SS,RS)-SAmE is subjected to lyophilization.
3. Process according to claim 1 or 2, wherein the pH value in step (a) is 1-2.
- 30 4. Process according to any of the preceding claims, wherein the preparation of the lysate in step (b) takes place by passing the yeast through a cell-breaking equipment.
5. Process according to any of the preceding claims,
35 wherein the temperature is 2-5°C.

- 13 -

6. Process according to any of the preceding claims, wherein the enriched yeast contains at least 8-10 g/l of (SS,RS)-S-adenosyl-L-methionine.

7. Process according to any of the preceding claims,
5 wherein the pharmaceutically acceptable acid is selected from sulphuric acid and paratoluensulphonic acid.

8. A pharmaceutically acceptable salt of (SS,RS)-S-adenosyl-L-methionine obtainable by the process according to any of the preceding claims.

INTERNATIONAL SEARCH REPORT

Application No

PCT/EP 01/03633

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07H19/16 C12P19/40 C07H1/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07H C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 621 056 A (GENNARI FEDERICO) 4 November 1986 (1986-11-04) the whole document	1-8
X	J. L. HOFFMAN: "Chromatographic analysis of the chiral and covalent instability of S-adenosyl-L-methionine" BIOCHEMISTRY, vol. 25, 1986, pages 4444-4449, XP002154223 page 4445, left-hand column, paragraph 6 -right-hand column, paragraph 2	8
A	FR 2 531 714 A (NIPPON ZEON CO) 17 February 1984 (1984-02-17) cited in the application page 15, claims 1-4	1-8
	—/—	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *G* document member of the same patent family

Date of the actual completion of the international search

25 September 2001

Date of mailing of the international search report

04/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fac. (+31-70) 340-3016

Authorized officer

de Nooy, A

INTERNATIONAL SEARCH REPORT

tr International Application No

PCT/EP 01/03633

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3 954 726 A (FIECCHI ALBERTO) 4 May 1976 (1976-05-04) abstract	7

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Application No

PCT/EP 01/03633

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4621056	A	04-11-1986	IT 1169773 B 03-06-1987
			AT 31550 T 15-01-1988
			AU 569107 B2 21-01-1988
			AU 3232884 A 28-02-1985
			CA 1223535 A1 30-06-1987
			DE 3468234 D1 04-02-1988
			DK 402884 A ,B, 25-02-1985
			EP 0141914 A1 22-05-1985
			ES 535378 D0 16-06-1985
			ES 8506044 A1 16-10-1985
			FI 843315 A ,B, 25-02-1985
			JP 1736937 C 26-02-1993
			JP 4021478 B 10-04-1992
			JP 60070097 A 20-04-1985
			MX 7625 E 27-03-1990
			NO 843366 A ,B, 25-02-1985
FR 2531714	A	17-02-1984	JP 1532553 C 24-11-1989
			JP 59029700 A 16-02-1984
			JP 63005039 B 01-02-1988
			CH 653341 A5 31-12-1985
			DE 3329218 A1 01-03-1984
			FR 2531714 A1 17-02-1984
			IT 1169764 B 03-06-1987
US 3954726	A	04-05-1976	IT 1022036 B 20-03-1978
			LU 79038 A1 26-06-1978
			AR 202219 A1 23-05-1975
			AT 347604 B 10-01-1979
			AT 532574 A 15-05-1978
			AU 7068974 A 08-01-1976
			BE 816915 A1 16-10-1974
			CA 1024088 A1 10-01-1978
			CH 592686 A5 15-11-1977
			DD 112451 A5 12-04-1975
			DE 2430999 A1 16-01-1975
			DK 342974 A ,B, 17-03-1975
			ES 427674 A1 01-12-1976
			FI 195474 A ,B, 28-12-1974
			FR 2234891 A1 24-01-1975
			GB 1443587 A 21-07-1976
			HK 31680 A 20-06-1980
			HU 170262 B 28-05-1977
			IE 39517 B1 25-10-1978
			IL 45161 A 31-08-1978
			JP 905944 C 18-04-1978
			JP 50076215 A 21-06-1975
			JP 52035727 B 10-09-1977
			NL 7408688 A ,B, 31-12-1974
			NO 742320 A ,B, 20-01-1975
			RO 64135 A1 15-11-1978
			SE 408712 B 02-07-1979
			SE 7408388 A 30-12-1974
			SU 576922 A3 15-10-1977
			US 4028183 A 07-06-1977
			YU 180174 A1 31-08-1982
			CS 183744 B2 31-07-1978
			RO 63045 A1 15-06-1978

INTERNATIONAL SEARCH REPORT

information on patent family members

In International Application No

PCT/EP 01/03633

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 3954726	A	ZA 7404161 A	25-06-1975